

Statistics of the single mode light in the $\chi^{(3)}$ transparent medium

V.N.Gorbachev*and A.I.Zhiliba†

February 1, 2008

Department of Physics, University of Sankt-Petersburg,
Department of Physics, Tver State University*,
170006,Tver, Private Bag 0609†, Russia

Abstract

The quantum statistics of the light in the transparent medium with cubic nonlinearity is considered. Two types of transparent media are treated,namely, the cold transparent medium with a ground working level and the inversion-free medium with the lasing levels of the same population. The spectra of light fluctuation are found on the basis of both Scully-Lamb and Haken theories. The conditions for the use of effective Hamiltonian are determined. Basing on the exact solution of the Fokker-Plank equation for the Glauber-Sudarshan P-function the inversion-free medium with cubic nonlinearity is shown to be the source of spontaneous radiation with non-Gaussian statistics.

Introduction

Four-wave-mixing processes (FWM) are of particular interest for the study of nonclassical states of the light as predicted by Yuen and Shapiro [1] and observed by Slusher et al. [2]. The quantum theory of FWM is based on two general approaches. The first is the Effective Hamiltonian Approach (EHA) when a nonlinear medium is described by a c-number coupling constant in an effective Hamiltonian of interaction. The starting point of the second approach is the master equation for the field obtained by adiabatic elimination of the atomic variables. EHA has been used to consider the time evolution of the number of optical systems [3]. Loudon et al have developed the quantum theory of propagation of the light and examined squeezing produced by self-phase modulation in fiber and in the nonlinear Saqnac interferometer with the help

*email: vn@VG3025.spb.edu

†email: Anatoly.Zhiliba@tversu.ru

of the effective Hamiltonian [4]. The effective Hamiltonians are usually introduced phenomenologically and occasionally exact solution may be obtained. EHA seems to be attractive because it requires only susceptibilities to describe the nonlinear medium. For three-photon parametric phenomena the effective Hamiltonian can be obtained if the $\chi^{(2)}$ medium is assumed to be transparent as it has been shown in [6]. As to the case of $\chi^{(3)}$ the above mentioned conditions [6] for the use of effective Hamiltonian construction would not be expected due to the existence of real transitions. It is well known that the fluctuation-dissipation theorem (FDTh) does not exist in the case of four-index susceptibilities of a $\chi^{(3)}$ medium [5].

The second approach has been developed for two-level models: the Scully-Lamb model (SLM) for the excited-state atom [7] and the Haken model (HM) for the ground-lower level atom [8],[9]. Walls and Reid have shown that statistics of the light of nondegenerate FWM under linear theory over two weak modes is the same in both HM and SLM [9]. For the single-mode FWM the master equations of HM and SLM are markedly different and the two models are in agreement only for the particular case of the linear theory [6].

The difference between HM and SLM arises not only from the type of the two-level atom. Formalism of the Haken theory is based on the multi-atom problem [10]. In contrast the starting points of the Scully-Lamb theory is the single-atom problem [11]. In this view these theories are sometimes classified as the multi-atom and single-atom approaches [12]. The problem is to find the most suitable approach [12], [13]. We believe that the HM and SLM describe an optical system in a quite different manner. It is an optical cavity with atoms inside for the HM while in the case of the SLM the atomic beam is injected into a cavity like in a maser or micromaser. It is clear that statistics of the light produced by the two systems may be substantially different. Here we use terms HM and LSM to take into account both the type of the approach (multi- or single-atom) and the type of the two-level atom. In the present paper we are aimed to apply our previous calculation within the framework of the Lax-Louisell model (LLM) to consider whether the effective Hamiltonians may be obtained for the $\chi^{(3)}$ medium and whether the statistics of the light is different for HM and LSM.

We discuss in this paper an optical scheme including the cavity with nonlinear medium inside and the classical injected signal. The $\chi^{(3)}$ medium is described by the Lax-Louisell model which may be directly reduced to the HM and SLM. Two kinds of transparent medium are considered. In Section 1 spectra of fluctuation of the output light is calculated for the cold-transparent medium for which only the lower working level is populated and the frequency of the light is far from the atomic resonance. In Section 2 we consider the inversion-free medium when the population of both working levels are equal. The semi-classical Maxwell theory predicts no evolution of the light in the inversion-free medium. From this point of view the medium is transparent. Nevertheless it is a source of the light analogous with that of spontaneous radiation. With the use of the exact solution of the Fokker-Planck equation for the Glauber-Sudarshan P-function the statistics of the light is found to be non-Gaussian.

1 The cold transparent medium

1.1 Initial equations

Let us consider a system consisting of N the similar three-level fixed atoms shown in Fig.1a. It is the Lax-Louisell model when the incoherent pumping provides the populations f_1^s and f_2^s of working levels 1 and 2, respectively, under conditions of the field absence. The medium are placed into the high Q cavity (see Fig.1b) at ω frequency, which can differ from the injected field frequency ω_L . The injected signal is considered to be monochromatic coherent wave.




Fig. 1a.




Fig.1b.

On the base of this model the master equation was derived by the adiabatic elimination of atomic variables in [6] where the field interaction with working transition was described by the Hamiltonian written in the dipole and rotating-field approximation. The resulted equation was the Fokker-Planck one for the

Glauber-Sudarshan quasi-probability.

$$\partial_t P = (\partial_\alpha \mu_1 + \partial_{\alpha\alpha}^2 \mu_2 + \partial_{\alpha\alpha}^2 \mu_3 + c.c.)P + R'_F P, \quad (1)$$

where the coefficients μ_i (correct to g^4) for the medium with cubic nonlinearity are:

$$\begin{aligned} \mu_1 &= N \frac{g^2 \delta^*}{|\delta|^2} \alpha (1 - \beta |\alpha|^2) (f_1^s - f_2^s) \\ \mu_2 &= -N \alpha^2 \frac{g^2 \delta^*}{|\delta|^2} \left\{ \frac{g^2 \delta^*}{|\delta|^2} (f_2^s - f_1^s) \left(\frac{1 - f_1^s}{\Gamma_1} + \frac{f_2^s}{\Gamma_2} \right) \right. \\ &\quad \left. + \frac{\delta^*}{|\delta|^2} (f_2^s - f_1^s) \right\} + \frac{\beta \delta}{2\gamma} f_2^s \\ \mu_3 &= -N \frac{g^2 \delta^*}{|\delta|^2} \left\{ \frac{g^2 |\alpha|^2}{|\delta|^2} (f_2^s - f_1^s) \left\{ \frac{\delta}{\Gamma_2} f_2^s + \frac{\delta}{\Gamma_1} (1 - f_1^s) \right. \right. \\ &\quad \left. \left. + f_2^s - f_1^s \right\} + \beta |\alpha|^2 \frac{\Gamma_1}{\Gamma_1 + \Gamma_2} (f_2^s - f_1^s) + \beta |\alpha|^2 \frac{\delta^*}{2\gamma} f_2^s - f_2^s \right\} \\ \delta &= \gamma + i\Delta \\ \Delta &= \omega_{21} - \omega \end{aligned} \quad (2)$$

γ and ω_{21} are the constant of the transverse relaxation and the frequency of the atomic transition respectively, $g = d\hbar^{-1}(\hbar\omega/2V\varepsilon_0)^{1/2}$ is the coupling constant, d is the dipole moment of the working transition, and $\beta = 2g^2\gamma(\Gamma_1 + \Gamma_2)(\Gamma_1\Gamma_2|\delta|^2)^{-1}$ is the saturation parameter. The relaxation constants Γ_1 and Γ_2 are formed from the set of initial constants:

$$\begin{aligned} \Gamma_2 &= (\gamma_{\downarrow} + \gamma'_2) + (\gamma_{\uparrow} + \gamma'_1)(\Lambda_1 + \Lambda_2 + \gamma_2)(\Lambda_1 + \Lambda_2 + \gamma_1)^{-1} \\ \gamma'_k &= \gamma_k(1 - \Lambda_k(\Lambda_1 + \Lambda_2)^{-1}) \\ k &= 1, 2. \end{aligned}$$

The constant Γ_1 can be obtained from Γ_2 by replacing $\gamma_{\uparrow} \rightarrow \gamma_{\downarrow}$, $\gamma_{\downarrow} \rightarrow \gamma_{\uparrow}$, $2 \rightarrow 1$, $1 \rightarrow 2$. The term

$$R'_F = \partial_\alpha \left(\frac{1}{2} C \alpha - a_0 \exp(i(\omega - \omega_L)t) \right) + c.c.$$

simulates the input and output of radiation from the cavity [11],[13]. Here $C = C_{in} + C_{out}$, $C_{in(out)} = cT^{in(out)}L^{-1}$, c is the light velocity, $T^{in(out)}$ is the transmission coefficient of the input and output mirrors, L is the cavity perimeter, and a_0 is the amplitude of the injected field in inside the cavity. This amplitude is considered to be real.

Equation (1) describes the linear and nonlinear processes of interaction. One can introduce the macroscopic susceptibilities for these processes.

$$\begin{aligned} \chi^{(1)}(\omega) &= N g^2 (\omega_{21} - \omega - i\gamma)^{-1} \\ \chi^{(3)}(\omega) &= \beta \chi^{(1)}(\omega) \end{aligned} \quad (3)$$

The Maxwell equation for the complex amplitude of the field inside the cavity $z = \int d^2\alpha P(\alpha, t)$, follows from Eq.(1) Accounting for Eq.(3) this equation can be written as

$$\partial_t z = ((i\kappa_1 - \kappa_2)z + (-i\chi_1 + \chi_2)z|z|^2)(f_1^s - f_2^s) - \frac{C}{2}z + a_0 \exp(i(\omega - \omega_L)t) \quad (4)$$

where the real and imaginary parts of the susceptibilities are separated out

$$\begin{aligned} \chi^{(1)} &= \kappa_1 + i\kappa_2 \\ \chi^{(3)} &= \chi_1 + i\chi_2 \end{aligned}$$

Since Eq. (1) is written correct up to g^4 , the dimensionless intensity of the field cannot be large.

$$\beta|z|^2, \quad \beta\langle|\alpha|^2\rangle \ll 1 \quad (5)$$

We consider the cold transparent medium assuming detuning $x = \Delta/\gamma$ to be large.

$$|x| \gg 1 \quad (6)$$

Under these conditions the real and imaginary parts of the susceptibilities are of essentially different order.

$$\begin{aligned} \kappa_1 &\sim Ng^2/\Delta, \quad \chi_1 \sim Nd^4/\Delta^3 \\ \kappa_2 &= \kappa_1/x, \quad \chi_1 = \chi_1/x \end{aligned}$$

Therefore the linear and nonlinear dispersion described by the susceptibilities κ_1 and χ_1 have a dominant role in a transparent medium where $f_2^s = 0$. The processes of absorption can be neglected setting $\kappa_1, \chi_2 = 0$. However the last-mentioned fact means that the main losses of the field in the cavity are connected only to the radiation output, i.e., $\kappa_2, \chi_2|z|^2 \ll C/2$.

As a result for the cold ($f_2^s = 0$) transparent medium we have:

$$\begin{aligned} \mu_1 &= [-i\kappa_1\alpha + i\chi_1\alpha|\alpha|^2]f_1^s \\ \mu_2 &= \alpha^2\chi_1(1 + \Gamma_1/\Gamma_2)^{-1}f_1^s(-\frac{x}{2}(1 - f_1^s) + \\ &\quad + i(f_1^s - 1 - f_1^s\Gamma_1/2\gamma)) \\ \mu_3 &= |\alpha|^2\chi_1(1 + \Gamma_1/\Gamma_2)^{-1}f_1^s(1 - f_1^s)\frac{x}{2} \end{aligned} \quad (7)$$

The Lax-Louisell model allows a direct transition to HM and SLM. In [6] the necessary conditions were formulated which can be reduced to the set of inequalities between constants $\gamma_\uparrow, \gamma_\downarrow$ and γ'_1, γ'_2 .

So, putting in Eq.(7)

$$\begin{aligned} \Gamma_1 &= \Gamma_2 = \gamma_\uparrow + \gamma_\downarrow = \gamma_\parallel, \quad f_1^s + f_2^s = 1 \\ \gamma &= \gamma_\perp \\ \gamma_\parallel/2\gamma_\perp &= f_c \end{aligned}$$

we obtain the Fokker-Planck equations for HM.

$$\partial_t P = (\partial_\alpha(-i\kappa_1\alpha + i\chi_1\alpha|\alpha|^2) + \partial_{\alpha\alpha}^2(-\frac{i}{2}\chi_1\alpha^2 f_c) + c.c.)P + R'_F P, \quad (8)$$

In this equation the various types of relaxation are described $0 < f_c \leq 1$, the f_c characterizes the difference between longitudinal and transversal relaxation velocities. The case when $f_c = 1$ describes the radiation decay only. We note that Eq.(8) agrees with equation obtained in [14].

Putting in Eq.(7)

$$\begin{aligned} \Gamma_k &= \gamma_k \\ (f_k^s)^2 &= 0, \quad f_k^s \neq 0, \quad k = 1, 2 \end{aligned} \quad (9)$$

we obtain the SLM equation.

$$\begin{aligned} \partial_t P &= f_1^s(\partial_\alpha(-i\kappa\alpha + i\chi\alpha|\alpha|^2) \\ &+ \partial_{\alpha\alpha}^2(-\chi_1(1 + \gamma_1/\gamma_2)^{-1}\alpha^2(i - x/2)) \\ &+ \partial_{\alpha\alpha}^2(\chi_1(1 + \gamma_1/\gamma_2)^{-1}|\alpha|^2 x/2) + c.c.)P + R'_F P \end{aligned} \quad (10)$$

The conditions (9) include the requirement for weak pumping $(f_k^s)^2 = 0, f_k^s \neq 0$, when the terms proportional to $(f_k^s)^2$ are absent. Formally, it is the consequence of the single-atom approach of the SLM. Physically, SLM describes optical systems in which the atoms are injected into the cavity being in a prepared state and rapidly escaped from the interaction zone. The atoms output from the cavity is modeled by their output from the working levels. On one hand, it requires an introduction of two-level systems with excited levels, and, on the other hand, it allows the use of the single-atom approach.

The Lax-Louisell model discussed here a priori describes the atom with two excited working levels. As it follows from Eq.(7) at

$$f_1^s = 1, \quad \Gamma_1\Gamma_2((\Gamma_1 + \Gamma_2)\gamma)^{-1} = f_c \quad (11)$$

the Fokker-Planck equations in the Haken and the Lax-Louisell models are the same. It means that the statistics of the field formed by two-level atoms with ground state and that of the field formed by atoms with two excited levels is the same. Therefore under conditions (11) the differences between LSM and LLM mainly depend upon the type of physical system rather than the type of two-level atoms. We note that these differences disappear when the linear over interaction theory is used. This theory is used as an example in the case of the linear medium or in the case of multi-photon interaction described by the effective Hamiltonian of the field in the linear approximation [14].

1.2 The effective Hamiltonian

The effective Hamiltonian introduced phenomenologically is often used to describe the four-wave mixing in the transparent medium with cubic nonlinearity.

The Hamiltonian is

$$H_{eff} = \frac{1}{2}\hbar k a^{+2} a^2, \quad (12)$$

where k is the coupling constant. The exact equation for the Glauber-Sudarshan quasi-probability written on the base of Eq.(12) has the Fokker-Plank equation form

$$\partial_t P = (\partial_\alpha (ik\alpha|\alpha|^2) + \partial_{\alpha\alpha}^2 (-\frac{i}{2}k\alpha^2) + c.c.)P.$$

For the cold transparent medium described by the Lax-Louisell model the effective Hamiltonian appears solely in the case when

$$\begin{aligned} f_1^s &= 1, \\ \Gamma_1 \Gamma_2 ((\Gamma_1 + \Gamma_2)\gamma)^{-1} &= 1. \end{aligned} \quad (13)$$

In the HM it means that $f_c = 1$, i.e., the decay is of the purely radiation character. Under conditions of Eq.(13) the equation for the electromagnetic field density matrix ρ following from Eq.(7) or Eq.(8) takes the form

$$\begin{aligned} \partial_t \rho &= -i\hbar^{-1}[H_1 + H_2, \rho] + R_F \rho \\ H_1 &= -\hbar \kappa_1 a^+ a \\ H_2 &= \frac{1}{2}\hbar \chi_1 a^{+2} a^2 \end{aligned} \quad (14)$$

where the operator $R_F \rho$ in P -representation corresponds to the $R'_F \rho$ operator. In contrast to the effective Hamiltonian approach (EHA) based on Eq.(12) the effect of the linear dispersion described by the Hamiltonian H_1 is taken into account in Eq.14.

Physically speaking the linear dispersion tends to the change of mode frequency $\omega \rightarrow \omega - \kappa_1$. It results the additional detuning between the injected field frequency and the cavity mode. The role of this factor was discussed in [9] and, as will be shown, it substantially affects the fluctuation dynamics and the noises. If the injected signal frequency is such that detuning is absent, i.e., $\omega_L - (\omega - \kappa_1) = 0$, the linear dispersion is not important in the transparent medium. In such a case the description by EHA with $k = Re\chi^{(3)}$ coincides with that by Eq.(14). The effective Hamiltonian principally does not arise in the SLM due of the terms proportional to x .

1.3 The Langevin equations

The condition of the small fluctuations allowing the linearization of the Fokker-Planck equation (FPE) is used as a basic approximation for the calculation of observed values. Basing on the linearized equation we write the Langevin equations which are convenient to obtained the fluctuation spectrum. We introduce polar coordinates

$$\alpha\alpha^* = I, \quad \varphi = (1/2i)\ln(\alpha/\alpha^*)$$

The fluctuations of intensity and phase around its semi-classical values are assumed to be small

$$\begin{aligned} I &= U + \varepsilon & \varepsilon &\ll U \\ \varphi &= \varphi_0 + \psi & \psi &\ll \varphi_0 \end{aligned} \quad (15)$$

where $U = |z|^2$ and $\varphi_0 = \arg z$ are the steady-state solutions of the semi-classical equations (4), which for transparent medium take the form :

$$\begin{aligned} \partial_t U &= -CU + 2a_0\sqrt{U} \cos \varphi_0 = 0 \\ \partial_t \varphi &= \kappa_1 - \chi_1 U + a_0 \sin \varphi_0 / \sqrt{U} = 0 \end{aligned} \quad (16)$$

In Eqs.(16) we assumed the phase of the injected field to be equal to zero and $\omega_L = \omega$. As a result we obtain the Langevin equation

$$\partial_t \begin{pmatrix} \varepsilon \\ \psi \end{pmatrix} = \begin{pmatrix} -A & A_{12} \\ A_{21} & -A \end{pmatrix} \begin{pmatrix} \varepsilon \\ \psi \end{pmatrix} + D^{1/2} \eta(t) \quad (17)$$

where $\eta(t)$ is the two-dimensional Winer process and the elements of the drift A and diffusion D matrix for the three considered models are listed in Table 1. Note that the linear dispersion affects only the fluctuations dynamics, and in SLM the source of the phase noise appears.

Table 1.

	EHA	HM	SLM
A	C/2		
A_{12}	$2kU^2$	$2\chi_1 U^2 - 2U\kappa_1$	
A_{21}	$-\frac{3}{2}k$	$-\frac{3}{2}\chi_1 + \frac{1}{2}\kappa_1/U$	
$D_{\varepsilon\phi}$	$-kU$	$-\chi_1 U f_c$	$-\chi_1 U$
$D_{\psi\psi}$	0		$\frac{1}{2}x\chi_1$

The phase and the intensity fluctuation are coupled dynamically $A_{12}, A_{21} \neq 0$ and statistically $D_{\varepsilon\psi} \neq 0$. This fact leads to two circumstances. First, both the intensity and phase fluctuations is produced by two sources $D_{\varepsilon\psi}$ and $D_{\psi\psi}$. Second, under steady-state the fluctuations being attenuated oscillate with the frequency Ω . To obtain Ω we write closed equations for the averaged values. These equations follow from Eq.(17).

$$\begin{aligned} (\partial_t^2 + 2A\partial_t + A^2 + \Omega^2)\sigma &= 0 \\ \sigma &= \langle \varepsilon \rangle, \langle \psi \rangle \\ \Omega^2 &= -A_{12}A_{21} \end{aligned} \quad (18)$$

In HM and SLM

$$\Omega^2 = \kappa^2(1 - \beta U)(1 - 3\beta U),$$

where the dimensionless intensity $\beta U \ll 1$ since the initial equation (1) is correct up to g^4 . Let

$$\beta U \ll 1/3 \quad (19)$$

Then $\Omega^2 = \kappa^2$ and Ω takes a new meaning of the damped harmonic oscillator frequency described by Eq.(18). As a result under steady-state the fluctuations of phase and intensity damp with velocity $C/2$ and oscillate at the frequency Ω .

In HM and SLM under conditions (19) the dynamics of the fluctuations are defined solely by the linear dispersion.

$$A_{12} = -2U\kappa_1, \quad A_{21} = \frac{1}{2}\kappa_1/U$$

where the expression for U follows from Eqs.(16) under conditions (19).

$$U = a_0^2(A^2 + \kappa_1^2)^{-1} \quad (20)$$

The output power $P = \hbar\omega cUT^{out}L^{-1}$ will be connected to the power of the injected light P_0 with the use of Eq.(20) as

$$P = P_0 \frac{1}{1+t^2} \cdot \frac{4\epsilon}{(1+\epsilon)^2}$$

where

$$t = 2\kappa_1/C, \quad \epsilon = T^{in}/T^{out} \quad (21)$$

1.4 The fluctuations spectra

For the homodyne balanced detection, when the signal is mixed with the field of local oscillator, the normal-ordered correlation functions of quadrature operator

$$X(t, \Theta) = a^+(t) \exp(i\Theta) + h.c. \quad (22)$$

are measured. The spectrum of the photocurrent $i^{(2)}(\omega)$ or the noise power spectrum can be presented in the form:

$$\begin{aligned} i^{(2)}(\omega) &= 1 + \eta g(\omega, \Theta) \\ g(\omega, \Theta) &= C^{out} \int_{-\infty}^{+\infty} d\tau e^{i\omega\tau} \langle : X(t, \Theta) X(t + \tau, \Theta) : \rangle \end{aligned} \quad (23)$$

where η is the quantum efficiency of the recording scheme, the points denote the normal ordering, and scale is determined so that the unit corresponds to the shot noise level. The value $g(\omega, \Theta)$ can be calculated by introduction of the Langevin variable

$$X(t, \Theta) = \alpha^*(t) \exp(i\Theta) + c.c. \quad (24)$$

which is c -number analog of the operator (22). So, it is easy to show that

$$g(\omega, \Theta) = C^{out} \int_{-\infty}^{+\infty} d\tau e^{i\omega\tau} \langle X(t, \Theta) X(t + \tau, \Theta) \rangle \quad (25)$$

The Eq. (25) is true when the Langevin equations are written from the FPE for the Glauber P-function or, as an example, the generalized P-representation. With the use of the small fluctuations (15)

$$X(t, \Theta) = 2\sqrt{U} \cos \varphi_0 + 2\sqrt{U} \sin(\Theta - \varphi_0)\psi(t) + \frac{1}{\sqrt{U}} \cos(\Theta - \varphi_0)\varepsilon(t)$$

Therefore the case when the phase of the local oscillator field coincides with that of the signal, i.e. $\Theta = \varphi_0$, corresponds to the measurement of spectrum of the intensity fluctuations, while when $\Theta - \varphi_0 = \frac{\pi}{2}$ the phase fluctuations are measured. Solving the Langevin equation (17) we obtain

$$\begin{aligned} g(\omega, \Theta) = & 2C^{out}[(\omega^2 + S_1^2)(\omega^2 + S_2^2)]^{-1} \cdot \\ & \cdot (D_{\psi\psi}[(\mu A + \nu A_{12})^2 + \mu^2 \omega^2] \\ & + 2D_{\epsilon\psi}[(\mu A + \nu A_{12})(\mu A_{21} + \nu A) + \mu\nu\omega^2]) \end{aligned}$$

where $\mu = \sqrt{2U} \sin(\Theta - \varphi_0)$, $\nu = \frac{1}{\sqrt{2U}} \cos(\Theta - \varphi_0)$, and $S_{1,2}$ are the roots of the equation $(S + A)^2 - A_{12}A_{21} = 0$. Introducing the scaled frequency $\bar{\omega} = \omega/A$, $t = \kappa_1/A$ we represent the obtained spectrum in the form

$$g(\omega, \Theta) = \frac{4}{1 + \epsilon} \frac{W(\Theta - \varphi_0) \cdot t^2 + V(\Theta - \varphi_0) \cdot \bar{\omega}^2}{(\bar{\omega}^2 + 1 - t^2)^2 + 4t^2}$$

For the measurement of the intensity and the phase fluctuations the coefficients are presented in Table 2. For SLM the case $\gamma_1 = \gamma_2$, $f_1^s = 1$ is taken as before.

	EHA	HM	LSM
$W(0)$	-2/3	$2\beta U f_c$	$2\beta U(1 + \frac{\pi t}{2})$
$V(0)$	0		
$W(\frac{\pi}{2})$	2	$-2\beta U f_c$	$-2\beta U(1 - \frac{\pi}{2t})$
$V(\frac{\pi}{2})$	0		$\beta U x t$
t	$\sqrt{3kU}/A$	κ_1/A	

Table 2.

When EHM and HM are compared one can see that the effect of dispersion is diametrically opposite behavior both for the phase and amplitude fluctuations. However on the basis of both approaches compaction arises leading to a decrease in the shot noise by a factor of not more than 3. Indeed the noise reduction may be obtained in nonzero frequency range. This behavior is understood by suggestion that the dispersion rotates the squeezing ellipse.

To determine the degree of squeezing for the optical scheme under consideration we need to choose the optimal phase of the local oscillator $\Theta = \Theta_0$. To this end we represent $g(\omega, \Theta)$ in the form $g = A + B \sin 2\Theta + C \cos 2\Theta$ and determine the Θ_0 value from the condition $\tan 2\Theta_0 = B/C$. As a result the fluctuation spectrum takes a form

$$g(\omega, \Theta_0) = \frac{2}{1 + \epsilon} [(\bar{\omega}^2 + 1 - t^2)^2 + 4t^2]^{-1} G(\omega)$$

where

$$\begin{aligned}
G_{EHA}(\omega) &= \frac{2t}{3}(2t \pm (4t^2 + 3(1 - t^2 + \bar{\omega}^2)^2)^{1/2}) \\
G_{HM}(\omega) &= \pm 2f_c \beta u |t| \cdot M \\
G_{LSM}(\omega) &= 2\beta U x t [x(1 + \bar{\omega}^2 + t^2) \pm M] \\
M &= [(1 + t^2)^{-1}[(1 + \bar{\omega}^2 + t^2)^2 + t^2(1 + t^2 - \bar{\omega}^2)^2]]^{1/2}
\end{aligned}$$

In the zero-frequency range

$$\begin{aligned}
g_{EHA}(0, \Theta) &= \frac{4}{3} \frac{t}{(1 + t^2)^2} (2t \pm \sqrt{4t^2 + 3(t^2 - 1)^2}) \\
g_{HM}(0, \Theta) &= \pm 4f_c \beta U \frac{|t|}{|t|^2 + 1} \quad \text{squeezing} - 2/3 \\
g_{LSM}(0, \Theta) &= \pm \frac{4}{x} \beta U \frac{t}{t^2 + 1} \quad \text{weaksqueezing}
\end{aligned}$$

For EHM the noise can be suppressed not only in the zero-frequency range. Choosing the local oscillator field phase using the condition $\tan \Theta = (\sqrt{3}t)^{-1}$, we find

$$g_{EHA}(\omega) = -\frac{8\bar{\omega}^2}{C[(\bar{\omega}^2 + 1 - t^2)^2 + 4t^2]} \cdot \frac{t^2}{1 + 3t^2}.$$

At the frequency $\bar{\omega}_e^2 = 1 + t^2$ we find the minimum

$$g_{min}(\bar{\omega}_e) = -\frac{2t^2}{C(1 + 3t^2)}$$

The maximum drop is equal 2/3.

2 Inversion-free transparent media

2.1 Initial equations

Let the pumping produce equal populations of working levels $f_1^s = f_2^s = f_0$. As it seen from FPE (1), in such a medium there is no polarization proportional to inversion. In this sense the inversion-free medium is transparent. Nevertheless the field can evaluate in such a media, especially in nonlinear one. At the same time it is seen directly from Eq.(1) that the contribution of second derivatives formed from the terms of μ_2 and μ_3 remains depending solely on the population of upper level f_2^s so the field can nevertheless evolve in such media, especially in nonlinear ones. Therefore the field in a inversion-free medium has a spontaneous radiation nature.

However the Eq.(1) does not permit the correct consideration of the field in a inversion-free medium because of the procedure of its derivation. The reason is that the noncommutativity of differential operators is neglected when the second derivatives are obtained. It has been shown in [6] that the noncommutativity contributes to the first derivatives in FPE that correspond to the semiclassical Maxwell's equations. If $f_1^s \neq f_2^s$ this contributions can be neglected. However, for inversion-free medium the first derivatives are produced by the noncommutativity only that why it must be taken into account. Using the formalism from [6] it is possible to obtain the master equation for the field with due regard for noncommutativity of differential operators. The Glauber-Sudarshan quasi-probability equation correct to g^4 for LLM has the form

$$\begin{aligned} \partial_t P = & \frac{1}{2} f_0 \left(\frac{\partial}{\partial \alpha} \alpha \chi_2 [3 + x^2 - 2ix] - \frac{\partial^2}{\partial \alpha^2} \alpha^2 \chi_2^2 (1 + x^2) \right. \\ & + \frac{\partial^2}{\partial \alpha \partial \alpha^*} [2\kappa_2 + \chi_2 ((x^2 - 1)|\alpha|^2 - 1 + \frac{2f_0(\Gamma_2 - \Gamma_1) - \Gamma_2}{\Gamma_1 + \Gamma_2})] \\ & + \frac{\partial^3}{\partial \alpha^* \partial \alpha^2} \alpha \cdot 2\chi_2 [1 - f_0 \frac{\Gamma_2 - \Gamma_1}{\Gamma_1 + \Gamma_2} + ix \frac{(f_0 - 1)(\Gamma_2 - \Gamma_1)}{\Gamma_1 + \Gamma_2}] \\ & \left. + \frac{\partial^4}{\partial \alpha^2 \partial \alpha^{*2}} 2\chi_2 f_0 \frac{\Gamma_1}{\Gamma_1 + \Gamma_2} + c.c) P \right) \end{aligned} \quad (26)$$

The polarization proportional to cubic susceptibility χ_2 appearing in inversion-free medium leads to evolution of the complex field amplitude $\langle \alpha \rangle = \int d^2 \alpha \alpha P(\alpha, t)$. So, according to Eq.(26)

$$\partial_t \langle \alpha \rangle = -\frac{1}{2} f_0 [\chi_2 (3 + x^2) - 2i\chi_1] \langle \alpha \rangle$$

The field amplitude damping and its steady-state value equals zero because $\chi_2 = 0$. From the physical point of view this result is related to the diffusion of the spontaneous radiation phase. Equation (26) can be approximated by the following FPE

$$\begin{aligned} \partial_t P = & \frac{1}{2} f_0 \left(\frac{\partial}{\partial \alpha} \alpha \chi_2 [3 + x^2 - 2ix] - \frac{\partial^2}{\partial \alpha^2} \alpha^2 \kappa_2 (1 + x^2) \right. \\ & \left. + \frac{\partial^2}{\partial \alpha \partial \alpha^*} [2\kappa + \chi_2 ((x^2 - 1)|\alpha|^2 - \frac{3}{2})] + c.c) P + R'_F P \right) \end{aligned} \quad (27)$$

where we set $\Gamma_1 = \Gamma_2$ and add the term $R'_F P$ assuming that the medium is placed in the cavity with the injected signal on. The FPE obtained can be solved exactly.

2.2 The exact solutions for free-inversion transparent medium

Let us represent the FPE (27) in the form

$$\begin{aligned} \partial_t P = & -\partial_\alpha (k\alpha + a_0) + \partial_{\alpha\alpha}^2 \alpha^2 \Lambda_{\alpha\alpha} \\ & + \partial_{\alpha\alpha^*}^2 (b + |\alpha|^2 \Lambda_{\alpha\alpha^*}) + c.c) P \end{aligned} \quad (28)$$

where a_0 and b are the real constants. To obtain the solutions the corresponding Langevin equations are used

$$\begin{aligned}\partial_t \alpha &= k\alpha + a_0 + f_\alpha \\ \partial_t \alpha^* &= (k\alpha + a_0)^* + f_{\alpha^*}\end{aligned}\tag{29}$$

where nonzero correlation functions for random forces have the form

$$\begin{aligned}\langle f_\alpha(t) f_\alpha(t+\tau) \rangle &= 2\langle \alpha^2 \rangle \Lambda_{\alpha\alpha} \delta(\tau) \\ \langle f_{\alpha^*}(t) f_{\alpha^*}(t+\tau) \rangle &= 2\langle \alpha^{*2} \rangle \Lambda_{\alpha\alpha}^* \delta(\tau) \\ \langle f_\alpha(t) f_{\alpha^*}(t+\tau) \rangle &= 2(b + \langle |\alpha|^2 \rangle) \Lambda_{\alpha\alpha^*} \delta(\tau)\end{aligned}$$

The formulated Langevin approach based on the FPE for the Glauber-Sudarshan quasi-probability was clasified in [14] as a "naive" one, because, for instance, the variables α and α^* cannot be interpreted as independent ones. Here we do not discuss neither the interpretation problems nor the mathematical aspects of stochastic differential equation theory, but only notice the following. First, the Langevin approach is in full accordance with the FPE (27) as far as concerned the calculation of observed values and based on Itto interpretation of stochastic integral. Second, all the results obtained below can be readily reproduced using, for example, the generalized quasi-probability $P(\alpha, \alpha^+)$, where α and α^+ are independent variables owing to the simple correspondence

$$P(\alpha, \alpha^*) \longleftrightarrow P(\alpha, \alpha^+) \quad \alpha, \alpha^* \longleftrightarrow \alpha, \alpha^+$$

Solving Eq.(29) we find

$$\begin{aligned}\alpha(t) &= z(t) + \int_0^t ds e^{k(t-s)} f_\alpha(s) \\ z(t) &= \alpha(0) e^{kt} + \frac{a_0}{k} (e^{kt} - 1)\end{aligned}\tag{30}$$

The expression for steady-state of complex amplitude of the field in the cavity can be consequently obtained.

$$\begin{aligned}\langle \alpha \rangle &= -a_0/k \\ Re k &< 0\end{aligned}$$

Using Eq.(30) we find the correlation function

$$\begin{aligned}Y(\tau, \Theta) &= \langle X(t, \Theta) X(t+\tau, \Theta) \rangle, \quad \tau > 0 \\ X(t, \Theta) &= \alpha^* \exp(i\Theta) + c.c.\end{aligned}$$

To do this we need in average values of the type $\langle \alpha(t) \alpha(t+\tau) \rangle$, $\langle \alpha^*(t) \alpha(t+\tau) \rangle$ and so on. For example, using Eq.(30) we obtain:

$$\begin{aligned}\langle \alpha(t) \alpha(t+\tau) \rangle &= z(t) z(t+\tau) \\ &+ e^{kt} [\langle \alpha^2(t) \rangle - z^2(t) + e^{2kt} \int_0^t ds_1 \int_0^t ds_2 \\ &e^{-k(s_1+s_2)} \langle f_\alpha(s_1) f_\alpha(s_2) \rangle]\end{aligned}$$

Due to the delta-function properties this integral is equal to zero, consequently, at $t \rightarrow \infty$

$$\langle \alpha(t) \alpha(t + \tau) \rangle = \langle \alpha \rangle^2 + e^{k\tau} (\langle \alpha^2 \rangle - \langle \alpha \rangle^2)$$

As a result at $t \rightarrow \infty$ the expression for Y takes the form

$$\begin{aligned} Y(t, \Theta) = & \langle \alpha \rangle \langle \alpha^* \rangle + e^{-2i\Theta} \langle \alpha \rangle^2 \\ & + e^{kt} (\langle |\alpha|^2 \rangle - \langle \alpha \rangle \langle \alpha^* \rangle) \\ & + e^{2i\Theta} (\langle \alpha^2 \rangle - \langle \alpha \rangle^2) + c.c. \end{aligned} \quad (31)$$

The expression for the Fourier transformation is written as

$$\begin{aligned} Y(\omega, \Theta) = & 2Re \int_0^\infty d\tau e^{i\omega\tau} Y(\tau, \Theta) \\ Y(\omega, \Theta) = & \frac{4}{A} [(\bar{\omega}^2 + 1 - T^2)^2 + 4T^2]^{-1} \cdot \\ & \cdot ((S + Re\mu)(1 + \bar{\omega}^2 + T^2) + TIm\mu(\bar{\omega}^2 - 1 - T^2)) \end{aligned} \quad (32)$$

where

$$\begin{aligned} T = & Im k / A \\ \bar{\omega} = & \omega / A \\ A = & -Re k \\ S = & \langle |\alpha|^2 \rangle - \langle \alpha \rangle \langle \alpha^* \rangle \\ \mu = & \exp(-2i\Theta) (\langle \alpha^2 \rangle - \langle \alpha \rangle^2) \end{aligned}$$

It should be noted that Eq.(31) is valid for any diffusion coefficient of FPE (28). Therefore the main problem reduces to the determination of the steady-state variance S and μ .

Using Eq.(28) or the Langevin equations (29) the closed equations for S and μ can be obtained, which can be solved exactly. As a result, at

$$\begin{aligned} Re k & < 0 \\ Re(k + \Lambda_{\alpha\alpha^*}) & < 0 \\ Re(k + \Lambda_{\alpha\alpha}) & < 0 \end{aligned}$$

we obtain the necessary steady-state solutions:

$$\begin{aligned} \langle |\alpha|^2 \rangle &= \left(\frac{a_0^2}{|k|^2} (k + k^*) - 2b \right) (k + k^* + 2\Lambda_{\alpha\alpha^*})^{-1} \\ S &= -2 \left(\frac{a_0^2}{|k|^2} \Lambda_{\alpha\alpha^*} + b \right) (k + k^* + 2\Lambda_{\alpha\alpha^*})^{-1} \\ \mu &= -e^{2i\Theta} \frac{a_0^2}{|k|^2} \cdot \frac{\Lambda_{\alpha\alpha}}{k + \Lambda_{\alpha\alpha}} \end{aligned} \quad (33)$$

It is interesting to note the following fact. The Fokker-Planck equations (8) and (9) for the SLM and the HM models reduce to Eq.(28) if the nonlinear

dispersion is neglected. For example, assuming that $\kappa \gg \chi_1|\alpha|^2$, i.e., $\beta|\alpha|^2 \ll 1$ we obtain in Eq.(28)

$$\begin{aligned} k &= -C/2 + i\kappa_1 \\ \Lambda_{\alpha\alpha} &= (-\frac{i}{2}\chi_1 f_c + \chi_1 x/4)f_1^s \\ \Lambda_{\alpha\alpha^*} &= f_1^s x \chi_1/4 \\ b &= 0 \end{aligned}$$

where $f_1^s = 1$, $x = 0$ should be set for the Haken model and $x \neq 0$, $f_c = 1$ for the Scully-Lamb one. The saturation parameter β can be considered in transparent medium as a small one (in the sense of the conditions (6)). This permits the neglect of $\Lambda_{\alpha\alpha}$ and $\Lambda_{\alpha\alpha^*}$ values in denominators of Eq.(33) when calculating S and μ . This approximation properly implies a linearization of the FPE near the steady-state $\langle\alpha\rangle$ obtained according to Eq.(30). As a result the solutions obtained in Section 1 both for the Haken and the Scully-Lamb models can be found using the presented formulas.

2.3 Fluctuation spectra

The term $\frac{1}{2}f_0\frac{\partial}{\partial\alpha}\alpha\chi_2(-2ix)$ in FPE (27) corresponds to the dispersion leading to the change of cavity mode frequency if $\omega \rightarrow \omega' - f_0\chi_2 \cdot x$. However, if the injected signal frequency is assumed to be $\omega_L = \omega'$ the dispersion plays no role. For this case the Eqs.(33) take the form

$$\begin{aligned} \langle|\alpha|^2\rangle &= (q_0 + \frac{a_0^2}{|k|^2}(1 + \frac{1}{2}\beta q_0(3 + x^2)))R \\ S &= (q_0 - \frac{\beta a_0^2}{2|k|^2}q_0(1 - x^2))R \\ \mu &= -\frac{\beta a_0^2}{2|k|^2}q_0(1 + x^2)R \exp(-2i\Theta) \\ R &= (1 + 2\beta q_0)^{-1} \end{aligned} \tag{34}$$

where the assumption $q_0 = 2f_0\kappa_2/C$ is used and $1 - 3\beta/4 \approx 1$ Because the value $k = -\frac{C}{2}[1 + \frac{1}{2}\beta q_0(3 + x^2)]$ is real the expression for spectrum $Y(\omega, \Theta)$ obtained according to Eq.(32) takes the simple form

$$Y(\omega, \Theta) = \frac{8(S + Re\mu)}{C[1 + \frac{1}{2}\beta q_0(3 + x^2)](1 + \bar{\omega}^2)}$$

Both the maximum and minimum values of Y is obtained at $\Theta = \frac{\pi}{2}(\max)$

$$S + Re\mu = q_0(1 + x^2\beta a_0^2/k^2)(1 + 2\beta q_0)^{-1}$$

and $\Theta = 0 (\min)$

$$S + Re\mu = q_0(1 - \beta a_0^2/k^2)(1 + 2\beta q_0)^{-1}$$

. For the case $\Theta = 0$, which corresponds to the measurement of amplitude fluctuations, the value $S + Re\mu > 0$. It follows from the fact that initial equation is correct to the order of g^4 i.e., only the cubic nonlinearity is taken into account. Therefore it is necessary to consider the field as not very strong one, i.e.,

$$\beta a_0^2/k^2 = \beta \langle \alpha \rangle \langle \alpha^* \rangle < 1, \quad \beta < |\alpha|^2 < 1$$

As a result the noise is found to be greater than the shot one. Physically it is clear because the inversion-free medium is the source of spontaneous radiation, the statistics of which, however, can be non-Gaussian.

2.4 Generation of light by the inversion-free medium

Let the injected signal and detuning be equal to zero. In such a medium the field with average number of photons

$$\begin{aligned} \langle n \rangle &= q_0(1 + 2\beta q_0)^{-1} \\ q_0 &= 2f_0\kappa_2/C \end{aligned} \tag{35}$$

and the Lorentzian profile of spectral line with halfwidth

$$\Delta\nu = \frac{C}{2}(1 + \frac{3}{2}\beta q_0)$$

is generated. Notice that the threshold of the generation is absent, and dimensionless intensity is limited:

$$\beta \langle n \rangle \leq 1/2$$

Consider the statistics of the generated light. Using Eq.(27) the Mandel parameter can be obtained:

$$\xi = \langle n \rangle (1 - \beta \langle n \rangle) \frac{1 + 2\beta q_0}{1 + \frac{5}{2}\beta q_0}$$

For the linear inversion-free medium $\beta = 0$, $\xi = \langle n \rangle$ i.e., the field statistics is of the Gaussian type. By contrast in the case under consideration $\frac{2}{5}\langle n \rangle \leq \xi < \langle n \rangle$. It means that the function of the photons number distribution is narrower than the Gaussian one. Therefore the spontaneous radiation statistics of the non-linear inversion-free medium is of essentially non-Gaussian nature.

3 Conclusion

The presented study of quantum statistics of single mode light in transparent $\chi(3)$ medium allows us to compare the intensity and phase fluctuation spectra obtained on the basis of both Haken and Scully-Lamb models. The performed

calculations clearly show how the light statistics in transparent $\chi(3)$ medium depends on the model medium excitation. It is seen that the light in Scully-Lamb model is more noisy than in the Haken one. Physically, it is rather evidently since in the SLM the medium subsystem is open. The atoms are injected into the cavity and leave the interaction zone rapidly. As a result, the SLM the medium excitation has additional source of noise with relation to the Haken model. It is shown in this paper that there are no any cases when EHA is valid in framework of Scully-Lamb model. But for the cold transparent medium Haken model with the lower working level to be ground state the traditional effective Hamiltonian of interaction was derived in particular case of absence of the collision-induced phase decay of atoms.

In the interesting case of the inversion-free $\chi(3)$ medium presented study leads to description of quantum evolution of light based on reduction of the Fokker-Plank equation from [6] to equation (20) presented paper. The obtained FPA has an exact solution. The statistics of generated light in this case is nongaussian that is photon number distribution is narrower than in the Gaussian statistics. At the same time there is no squeezing. Is this state of generated light quantum because this state and its evolution appear only in the framework of quantum description? It necessary that further study of the problem should be performed.

References

- [1] H.P.Yuen, J.H.Shapiro, Opt.Lett.**4**(1979)334
- [2] R.E.Slusher, L.W.Holberg., B.Yurke, J.C.Merta, J.F.Valley, Phys.Rev.Lett.**55** (1979)2409
- [3] J.Perina, V.Perinova, C.Sibilia, M.Bertolotti,Opt. Commun.**49**285(1984). R.Tanas, S.Kielich,Opt.Communications**45**351(1985). A.Belinskii,Zh.Eksp.Teor.Fiz.**99**748(1991).
- [4] K.J.Blow, R.Loudon, S.J.D.Phoenix,J.Opt.Soc.Am.**8** 1750(1991); Phys.Rev.A **45**8064(1992).
- [5] R.A.Stratanovich, Zh.Eksp.Teor.Fiz. **58** (1970) 1612.
- [6] V.N.Gorbachev, A.I.Zhiliba,Quantum Opt.**5** 193 (1993).
- [7] M.Sargent, D.Holm, M.Zubairy,Phys.Rev.**A 31** 3112 (1985)
- [8] R.Bonifacio, M.Gronchi, L.A.Lugiato,Phys.Rev.A **18** 2226 (1978)
- [9] M.D.Reid, D.F.Walls Phys.Rev.A **34** 4929 (1986)
- [10] H.Haken, H.Risken, W.Weidlich,Zs.Phys.**206** 355 (1967)
- [11] M.Scully, W.E.Lamb,Phys.Rev.**159** 208 (1967)
- [12] L.A.Lugiato, F.Casagrande, L.Pizzuto,Phys.Rev.**A 26** 3338 (1982)
- [13] R.Schack, A.Sizmann, A.Schenzle,Phys.Rev.A **43** 6303 (1991)
- [14] C.W.Gardiner *Handbook on stochastic Methods* (Springer Verlag,New-York,1985)